

Stable Parabolic Bundles over Elliptic Surfaces and over Orbifold Riemann Surfaces

Christian Gantz¹, Brian Steer
Mathematical Institute, Oxford OX1 3LB, UK
(gantz@maths.ox.ac.uk)

February 5, 2008

Mathematics Subject Classification (1991) 14J27 32L07 14H60 14D20

1 Introduction

If $q : Y \rightarrow \Sigma$ is an elliptic surface (to be made precise) then the induced map of fundamental groups is an isomorphism if we consider Σ as an orbifold, [U], [Dol]. Hence, we obtain a correspondence of flat bundles (by bundles we always mean complex vector bundles). Donaldson showed that each stable degree zero bundle over Y or Σ admits a unique Hermitian-Yang-Mills (or anti-self-dual) connection. In fact, the obvious conditions on a bundle $E' \rightarrow Y$ to come from Σ , namely $c_1(E') \in q^*H^2(\Sigma)$ and $c_2(E') = 0$, imply that this H.Y.M. connection is flat. So, there is a correspondence of stable degree zero bundles over Y and Σ . The generalisation to all degrees has been shown by Bauer, [B], via a direct proof in algebraic geometry.

Donaldson's result has been extended by several authors to parabolic bundles, theorem 5.2. This and the use of extension results for flat bundles, theorem 5.4, gives an identification of stable parabolic degree zero bundles over (Σ, P) and (Y, P') , where P is a finite collection of generic points in Σ and $P' = q^{-1}(P)$, with representations of the fundamental groups of $\Sigma - P$ and $Y - P'$, respectively. These groups are again isomorphic and so, we have a correspondence of stable parabolic degree zero bundles over (Y, P')

¹The first author was fully supported by the *Rhodes Trust*, Oxford, and the *Alfried Krupp von Bohlen und Halbach – Stiftung*, Essen

and over (Σ, P) . This extends readily to all degrees by tensoring with a parabolic line bundle.

Another proof for genuine bundles of any degree, not using parabolic bundles at all, relies on the correspondence between stable bundles over $\Sigma(Y)$ and representations of the fundamental group of circle bundles (i.e. Seifert fibred spaces) over $\Sigma(Y)$, [FS, p 63], [BO]. These groups are also isomorphic, see the proof of proposition 6.1.

Combining Bauer's arguments with Kronheimer & Mrowka's description of the moduli spaces of stable parabolic bundles, we show that these are complex manifolds if we fix determinants and if they are pull backs from Σ . Finally, we consider smooth parabolic bundles and produce some details about the correspondence of stable ones.

After recalling elliptic surfaces and parabolic bundles in the following two sections we take one section to state the results. More details of our work can be found in [G].

Acknowledgements

We are grateful to P. Kronheimer, S. Bauer, M. Lübke, T. Peternell, C. Okonek, S. Agnihotri and R. Plantiko for helpful remarks.

2 Elliptic surfaces

Throughout, let $q : Y \rightarrow \Sigma$ be an elliptic surface, i.e. Y is a compact complex surface, Σ a compact Riemann surface and $q^{-1}(\sigma)$ an elliptic curve for generic, i.e. all but finitely many, $\sigma \in \Sigma$, c.f. [GH], [BPV]. We will assume that any non-generic fibre is either a rational curve of multiplicity one with one self-intersection (called singular fibre) or a multiple elliptic curve and furthermore, that there is at least one singular fibre. Moishezon shows that all elliptic surfaces are deformation equivalent to the ones we consider, [Mo].

Theorem 2.1 (Ue, Dolgachev) *If $q : Y \rightarrow \Sigma$ has singular fibres and if $U_0 \subseteq \Sigma$ is an open ball such that $\pi^{-1}(U_0)$ contains all singular fibres but no multiple elliptic curves then $\pi^{-1}(U_0)$ is simply connected.*

If $Y_\sigma := q^{-1}(\sigma)$ has multiplicity $m > 1$, let $\tilde{U}$ and B be open discs in $\mathcal{C}$, $\phi : B \rightarrow U$ a chart with $U \subseteq \Sigma$ and $\phi(0) = \sigma$ and construct a uniformization of U by

$$\tilde{U} \xrightarrow{z^m} B \xrightarrow{\phi} U$$

where $\langle \eta = e^{2\pi i/m} \rangle = \mathbb{Z}_m \subseteq \mathcal{C}$ acts on $\tilde{U}$ in the standard way. After this construction is done for all multiple points $\sigma \in \Sigma$ we think of Σ as an orbifold, cf. [FS]. Bauer points out that for all orbifold Riemann surfaces there exists an elliptic surface over it.

The natural gauge-theoretic objects on orbifolds are V-bundles: a (local, complex) rank r V-bundle $E|_U$ is isomorphic to

$$(\tilde{U} \times \mathcal{C}^r, \mathbb{Z}_m) \rightarrow (\tilde{U}, \mathbb{Z}_m) \quad \text{with}$$

$$\eta(\tilde{u}, z_1, \dots, z_r) = (\eta\tilde{u}, \eta^{a_1}z_1, \dots, \eta^{a_r}z_r)$$

for some isotropies $(a_1, \dots, a_r) \in \{0, \dots, m-1\}^r$.

Theorem 2.2 (Furuta & Steer, Seifert) *Smooth V-bundles over Σ are classified by rank, degree (which is rational) and isotropies.*

For any $y \in Y_\sigma$ we can choose coordinates (z_1, z_2) on $U' \ni y$ such that $\phi^{-1} \circ q(z_1, z_2) = z_2^m$. Hence we can lift $\phi^{-1} \circ q$ locally to a regular map

$$\sqrt[m]{\phi^{-1} \circ q} = z_2 : U' \rightarrow \tilde{U}$$

uniquely up to the action of $\mathbb{Z}_m$. In particular, q is a map of orbifolds.

A divisor on Σ can be represented by a finite sum $D = \sum_{i \in I} \sigma_i n_i / m_i$ where $n_i \in \mathbb{Z}$ and m_i the multiplicity of $\sigma_i \in \Sigma$. The vertical divisors on Y are precisely the pull backs of divisors on Σ . Hence, [NasS], the line bundles $\mathcal{O}(D')$ over Y with vertical divisor correspond to the line V-bundles over Σ .

As q is regular away from finitely many points theorem 2.1 implies that $q_* : \pi_1(Y) \rightarrow \pi_1^V(\Sigma)$ is an isomorphism. Here, the orbifold fundamental group $\pi_1^V(\Sigma)$ is an extension of $\pi_1(\Sigma)$ by a torsion group with unipotent elements corresponding to the multiple points. The isomorphism implies that the first betti number of Y is even and therefore Y is Kahler, by [Mi]. The correspondence between representations of the fundamental group and flat bundles on manifolds extends to orbifolds via the construction with simply connected coverings. Hence, the flat bundles over Y correspond to the flat V-bundles over Σ .

3 Parabolic bundles

Let $P := \{p_1, \dots, p_n\}$ be a collection of generic points in Σ and put $P' := \{P'_j := q^{-1}(p_j)\}_{j=1 \dots n} \subseteq Y$. A parabolic V-bundle E over the pair (Σ, P) is

a V-bundle $E \rightarrow \Sigma$ with proper filtrations

$$E|_{P_j} = E_{j,1} \supset E_{j,2} \supset \dots \supset E_{j,l_j} \neq 0$$

and weights

$$0 \leq \alpha_{j,1} < \alpha_{j,2} < \dots < \alpha_{j,l_j} < 1$$

for all $j = 1, \dots, n$. We call $\mu_{j,k} := \text{rank}(E_{j,k}/E_{j,k+1})$ the multiplicity of $\alpha_{j,k}$. Let α_j be the diagonal matrix of rank equal to $\text{rank } E$ and with entries $\alpha_{j,k}$ ($k = 1, \dots, l_j$) with multiplicities. Then put $\alpha(E) := \{\alpha_1, \dots, \alpha_n\}$. We will sometimes write $|E|$ for the underlying genuine V-bundle.

Correspondingly, one defines parabolic bundles $E' \rightarrow (Y, P')$ where the $\{E'_{j,k}\}$ are bundles over P'_j with weights $\{\alpha'_{j,k}\}$ of multiplicities $\{\mu'_{j,k}\}$, encoded in $\alpha'(E')$. If the bundle and the filtrations are holomorphic we speak of a holomorphic parabolic bundle. Thinking of vectors in $E|_P$ as having (the obvious) weights ($+\infty$ being the weight of zero vectors), a morphism of parabolic bundles is a bundle map not decreasing the weight of any vector. In a direct sum of parabolic bundles the weight of a vector is the minimum of the weights of its projections.

Let us fix a Kahler metric on Y with $(1,1)$ -form ω' . When we use ordinary bundle invariants and operators (like c_i , $\det$ or $\deg$) on parabolic bundles we mean to apply them to the underlying bundles. Then we have

$$\text{par } c_1(E) := c_1(E) + \sum_{j=1}^n \text{Tr}(\alpha_j) \text{PD}(p_j) \in H^2(\Sigma, \mathbb{R})$$

and similarly, $\text{par } c_1(E') \in H^2(Y, \mathbb{R})$. Let $\text{par } \deg E := \langle \text{par } c_1(E), \Sigma \rangle \in \mathbb{R}$ and $\text{par } \deg E' := \langle \text{par } c_1(E') \cup \omega', Y \rangle \in \mathbb{R}$. We also have (since $P \cdot P = 0$)

$$\text{par } c_2(E') := c_2(E') + 2 \sum_{j=1}^n \sum_{k=1}^{l_j} \alpha'_{j,k} \text{PD}(d'_{j,k}) + \frac{1}{2} \text{par } c_1^2(E') \in H^4(Y, \mathbb{R})$$

where $d'_{j,k} := \deg(E'_{j,k}/E'_{j,k+1})$.

Definition 3.1 *A holomorphic parabolic V-bundle $\mathcal{E} \rightarrow (\Sigma, P)$ is called **stable** if for all non-zero parabolic maps $\mathcal{F} \rightarrow \mathcal{E}$, injective over some point in Σ and with $\text{rank } \mathcal{F} < \text{rank } \mathcal{E}$ we have*

$$\deg \mathcal{F} / \text{rank } \mathcal{F} < \deg \mathcal{E} / \text{rank } \mathcal{E}.$$

Similarly for $\mathcal{E}' \rightarrow (Y, P')$.

4 Statement of results

Being a map of orbifolds, $q : Y \rightarrow \Sigma$ induces pull backs of holomorphic parabolic V-bundles over Σ to holomorphic parabolic bundles over Y . Our main result is:

Theorem 4.1 *Pulling back induces a correspondence between stable parabolic V-bundles $\mathcal{E} \rightarrow (\Sigma, P)$ and those stable parabolic bundles $\mathcal{E}' \rightarrow (Y, P')$ with $\text{par } c_2(\mathcal{E}') = 0$ and $\text{par } c_1(\mathcal{E}') \in q^* H^2(\Sigma, \mathbb{R})$.*

In particular:

Theorem 4.2 *Pulling back induces a bijection between stable V-bundles $\mathcal{E}$ over Σ and those stable bundles $\mathcal{E}'$ over Y with $c_2(\mathcal{E}') = 0$ and $\det \mathcal{E}' = \mathcal{O}(D')$ for a vertical divisor D' .*

Theorem 4.2 has been shown (under some assumptions on the Kahler metric of Y , [B, p 511]) by Bauer first, using algebraic geometry. Our goal is a differential-geometric proof of theorem 4.1 using the correspondence of stable parabolic bundles and parabolic H.Y.M. connections.

Let $\mathcal{E} \rightarrow (\Sigma, P)$ be a stable parabolic bundle, $\mathcal{E}' := q^* \mathcal{E}$, $\mathcal{T} := \text{Par End}_0 \mathcal{E}$ and $\mathcal{T}' := q^* \mathcal{T} = \text{Par End}_0 \mathcal{E}'$. The important thing to note is that $\mathcal{T}$ and $\mathcal{T}'$ are holomorphic (V-)bundles. We have the deformation complexes (of smooth sections)

$$\begin{array}{ccccc}
 \Omega_{\Sigma}^0(\mathcal{T}) & \xrightarrow{\bar{\partial}_{\mathcal{T}}} & \Omega_{\Sigma}^{0,1}(\mathcal{T}) & \xrightarrow{\bar{\partial}_{\mathcal{T}}} & 0 \\
 \downarrow & & \downarrow q^* & & \downarrow \\
 \Omega_Y^0(\mathcal{T}') & \xrightarrow{\bar{\partial}_{\mathcal{T}'}} & \Omega_Y^{0,1}(\mathcal{T}') & \xrightarrow{\bar{\partial}_{\mathcal{T}'}} & \Omega_Y^{0,2}(\mathcal{T}')
 \end{array}$$

Let $E :=_{C^\infty} \mathcal{E}$, $\mathcal{O}(D) = \det \mathcal{E}$ and $\mathcal{M}(E, D)$ be the space of stable structures on E with determinant $\mathcal{O}(D)$. Similarly, define $\mathcal{M}(E', D')$ for $E' := q^* E$ and $D' := q^* D$. Using the extension of standard deformation

theory, [DK], to parabolic bundles makes $\mathcal{M}(E', D')$ into a Hausdorff complex space, see [KM95, p 83] and [Mu], with a description near $\mathcal{E}'$ given by the zero set of a holomorphic map

$$H_{\bar{\partial}}^{0,1}(Y, T') \rightarrow H_{\bar{\partial}}^{0,2}(Y, T').$$

By the vanishing of the second cohomology over Σ , $\mathcal{M}(E, D)$ is even a complex manifold with a chart near $\mathcal{E}$ given by $H_{\bar{\partial}}^{0,1}(\Sigma, T)$.

Proposition 4.3 (Bauer) *Stability of $\mathcal{E}$ implies that*

$$q^* : H_{\bar{\partial}}^{0,1}(\Sigma, T) \rightarrow H_{\bar{\partial}}^{0,1}(Y, T')$$

is an isomorphism.

This proposition and theorem 4.1 imply:

Corollary 4.4 *If $\mathcal{M}_{\mathcal{E}}(E, D)$ is the connected component of $\mathcal{E}$, correspondingly for $\mathcal{E}'$, then*

$$q^* : \mathcal{M}_{\mathcal{E}}(E, D) \rightarrow \mathcal{M}_{\mathcal{E}'}(E', D')$$

is an isomorphism of complex manifolds.

Proposition 4.5 *If $E' \rightarrow (Y, P')$ is a rank r parabolic bundle satisfying $\text{par } c_2(E') = 0$, $\text{par } c_1(E') \in q^* H^2(\Sigma, \mathbb{R})$ and if the space $\mathcal{M}(E')$ of stable structures on E' is non-empty then*

- (i) *there exists a unique line V -bundle $L \rightarrow \Sigma$ with $q^* L = \det E'$. Also, E' is uniquely determined by L and it's weights α' ;*
- (ii) *we have*

$$\mathcal{M}(E') = \bigsqcup_a \mathcal{M}(L, r, a, \alpha')$$

where the union is over isotropies $a \in \mathbb{Z}_{m_1} \times \dots \times \mathbb{Z}_{m_k}$ ($k = \#\{\text{marked points on } \Sigma\}$) compatible with L and $\mathcal{M}(L, r, a, \alpha')$ is the space of stable structures on the unique rank r parabolic V -bundle over (Σ, P) with determinant L , isotropies a and weights α' .

5 Stable parabolic bundles

Definition 5.1 *A (parabolic) hermitian metric h' on a parabolic bundle $E' \rightarrow (Y, P')$ (similarly for $E \rightarrow (\Sigma, P)$) is a hermitian metric on $E'|_{Y-P'}$ such that*

$$\forall y \in P'_j \quad E'_{j,k}(y) = \{s(y) \mid s \in \Gamma_{loc}(E') \text{ s.t. } h'(s(-)) = O(d(P'_j, -)^{\alpha'_{j,k}})\}.$$

For a holomorphic parabolic bundle $\mathcal{E}'$, h' induces a Chern connection on $\mathcal{E}'|_{Y=P'}$. So, one can talk of H.Y.M.-connections. Around P'_j , a parabolic connection has holonomy conjugated to α'_j , [KM93], and can therefore not be extended over P'_j , in general. A parabolic connection is called reducible if the parabolic bundle decomposes together with the connection. The bridge between algebraic and differential geometry is

Theorem 5.2 ([MS], [SW], [Mu], [B93], [NasS]) *Any degree zero holomorphic parabolic bundle over a complex Kahler surface or orbifold of dimension one is stable if and only if it admits an irreducible H.Y.M. metric, unique up to isomorphism. Furthermore, an H.Y.M. connection has finite action.*

The primary result, for genuine bundles, is due to Donaldson; Munari's proof relies on Simpson's work. A H.Y.M. connection over Σ is obviously flat.

Lemma 5.3 *If $\mathcal{E}' \rightarrow (Y, P')$ has $\text{par } c_2(\mathcal{E}') = 0$ and $\text{par } c_1(\mathcal{E}') \in q^* H^2(\Sigma, \mathbb{R})$ then a H.Y.M. connection is necessarily flat.*

Proof: By [Mu, p 100] or [G] we obtain the second (in particular the existence of) and, from [DK], the third equality in

$$0 = 8\pi^2(\text{par } c_2(\mathcal{E}') - \frac{1}{2}\text{par } c_1^2(\mathcal{E}')) = \int_{Y-P'} \text{Tr}(F^2) = \|F^-\|^2 - \|F^+\|^2.$$

□

Theorem 5.4 (Munari, Biquard, Simpson) *A holomorphic bundle over $Y - P'$ ($\Sigma - P$) with a flat hermitian metric extends uniquely (up to isomorphism) to a holomorphic parabolic bundle over (Y, P) ((Σ, P)) such that the hermitian metric becomes a parabolic metric.*

Now we prove theorem 4.1:

Proof: At first, we treat the special case of degree zero bundles. If $\mathcal{E} \rightarrow (\Sigma, P)$ is stable and of degree zero, let A be the unique irreducible flat parabolic connection on $\mathcal{E}$. By the uniqueness in theorem 5.4, $A|_{\Sigma-P}$ is still irreducible. The regularity of $q|_{Y-P'}$ away from finitely many points and theorem 2.1 imply that $(q|_{Y-P'})_* : \pi_1(Y-P') \rightarrow \pi_1^V(\Sigma-P)$ is an isomorphism. Hence, if $\mathcal{E}' := q^*\mathcal{E}$ and $A' := q^*A$ then A' is an irreducible flat parabolic connection and hence $\mathcal{E}'$ is stable.

Conversely, if $\mathcal{E}' \rightarrow (Y, P')$ is stable and of degree zero, let A' be the unique parabolic H.Y.M. connection, which is flat by lemma 5.3 and $A'|_{Y-P'}$ is irreducible. We push forward $(\mathcal{E}', A')|_{Y-P'}$ to $\Sigma-P$. By theorem 5.4, this push forward extends uniquely to a holomorphic parabolic bundle with irreducible parabolic connection $(\mathcal{E}, A) \rightarrow (\Sigma, P)$. As $q^*(\mathcal{E}, A)$ and $(\mathcal{E}', A')$ are isomorphic over $Y-P'$ the uniqueness of theorem 5.4 implies that they are isomorphic over (Y, P') . We are finished with the degree zero case.

To show the general case, fix a generic point $p \in \Sigma-P$ and let $\Delta := \int_{q^{-1}(p)} \omega'$. For each $d \in \mathbb{R}$ let $[d]$ be it's integer part, let $[p]$ be the holomorphic line bundle with divisor p and define the holomorphic parabolic line bundle $\mathcal{L}_d \rightarrow (\Sigma, p)$ to be $\mathcal{L}_d := [p]^{[d]}$ with weight $d - [d]$ over p . Let $\mathcal{L}'_d := q^*\mathcal{L}_d$ be the parabolic pull back. We have $\text{par deg } \mathcal{L}_d = d$ and $\text{par deg } \mathcal{L}'_d = \Delta d$.

Tensoring any rank r stable parabolic bundle $\mathcal{E} \rightarrow (\Sigma, P)$ ($\mathcal{E}' \rightarrow (Y, P')$) of degree $-dr$ ($-\Delta dr$) with $\mathcal{L}_d$ ($\mathcal{L}'_d$) gives a stable parabolic bundle of degree zero over $(\Sigma, P \cup \{p\})$ ($(Y, P' \cup \{q^{-1}(p)\})$). Furthermore, we have $\text{par } c_2(\mathcal{E}' \otimes \mathcal{L}'_d) = 0$. So we are reduced to the degree zero case. $\square$

We use Bauer's arguments, [B, p 514], to prove proposition 4.3:

Proof: The sheaf $\mathcal{T}$ can be considered as a sheaf on Σ or on $|\Sigma|$, the underlying Riemann surface of the orbifold Σ , see the correspondence between V-bundles and parabolic bundles in section 5 of [FS]. As sheaves, $q_*\mathcal{T}' = \mathcal{T}$. The Leray spectral sequence induces an exact sequence, [BPV, p 10],

$$0 \rightarrow H^1(\Sigma, \mathcal{T}) \xrightarrow{q_*} H^1(Y, \mathcal{T}') \rightarrow H^0(\Sigma, q_{*1}\mathcal{T}') \rightarrow \dots$$

where $q_{*1}\mathcal{T}'$ is the first direct image sheaf of $\mathcal{T}'$. It suffices to see that $H^0(\Sigma, q_{*1}\mathcal{T}') = 0$.

Relative duality, [BPV, p 99], gives

$$q_{*1}\mathcal{T}' = q_*(\mathcal{T}'^* \otimes \mathcal{K}_Y \otimes q^*\mathcal{K}_{|\Sigma|}^*)^* = \mathcal{T} \otimes q_*(\mathcal{K}_Y \otimes q^*\mathcal{K}_{|\Sigma|}^*)^*$$

where $\mathcal{K}_{|\Sigma|}$ is the canonical bundle. By [BPV, p 98, 161-162],

$$\mathcal{K}_Y \otimes q^* \mathcal{K}_{|\Sigma|}^* = q^* q_{*1} \mathcal{O}_Y^* \otimes \mathcal{O}_Y \left(\sum (m_i - 1) Y_{\sigma_i} \right)$$

where the sum is over the singular points σ_i of Σ , $q_{*1} \mathcal{O}_Y$ is locally free of rank one since all the other sheaves in this identity are, and

$$\deg(q_{*1} \mathcal{O}_Y) = -\chi(\mathcal{O}_Y).$$

In particular, $\mathcal{K}_Y \cdot \mathcal{K}_Y = 0$. Hence, $\chi(Y) = 12\chi(\mathcal{O}_Y)$, [GH, p 472], which is equal to the positive number of singular fibres, cf. [U]. Now, $\mathcal{O}_Y((m_i - 1)Y_{\sigma_i}) = q^* \mathcal{O}_\Sigma(\frac{m_i - 1}{m_i} \sigma_i)$. (This is in fact a trivial sheaf over Σ .) We obtain

$$q_{*1} \mathcal{T}' = \mathcal{T} \otimes q_{*1} \mathcal{O}_Y \otimes \mathcal{O}_\Sigma \left(\sum \frac{1 - m_i}{m_i} \sigma_i \right)$$

and any non zero section of this induces a non zero map $\mathcal{E} \rightarrow \mathcal{E} \otimes \mathcal{L}$ for a negative line V-bundle $\mathcal{L}$. This is ruled out by stability of $\mathcal{E}$. $\square$

6 Smooth parabolic bundles

Proposition 6.1 *Two smooth line V-bundles over Σ are isomorphic if their pull backs to Y are isomorphic.*

There is an equivalence between smooth line V-bundles over Σ and $H_V^2(\Sigma, \mathbb{Z})$, [FS]. As we don't have a sufficient theory of V-cohomology however, our proof is not by showing injectivity of $H_V^2(\Sigma, \mathbb{Z}) \rightarrow H^2(Y, \mathbb{Z})$.

Proof: It suffices to show that a smooth line V-bundle $L \rightarrow \Sigma$ is trivial if $L' := q^* L \rightarrow Y$ is trivial. Let us write $q' : SL' \rightarrow SL$ for the induced map of circle bundles. We use the isomorphism $q_* : \pi_1(Y) \rightarrow \pi_1^V(\Sigma)$ and the commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & K' & \longrightarrow & \pi_1(SL') & \xrightarrow{\pi'_*} & \pi_1(Y) & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow q'_* & & \downarrow q_* & & \downarrow \\ 0 & \longrightarrow & K & \longrightarrow & \pi_1^V(SL) & \xrightarrow{\pi_*} & \pi_1^V(\Sigma) & \longrightarrow & 0 \end{array}$$

where $K = \langle k \rangle$, $K' = \langle k' \rangle$ for regular fibres k of SL and k' of SL' . The rows are exact because bundles are always regular maps. By the five-lemma, q'_* is an isomorphism if $(k' \mapsto k) : K' \rightarrow K$ is an isomorphism. Certainly, it is surjective. Assume there is a V -homotopy $H : [0, 1] \times [0, 1] \rightarrow SL$ with boundary k^n . Since q is regular away from finitely many points, there exists $H' : [0, 1] \times [0, 1] \rightarrow Y$ lifting $\pi \circ H$. Hence there exists $\tilde{H} : [0, 1] \times [0, 1] \rightarrow SL'$ lifting H and H' . If $k \in S(L_x)$ then $\partial H' = \pi' \circ \partial \tilde{H} \subseteq Y_x$ and $\partial H = q' \circ \partial \tilde{H} = k^n$.

W.l.o.g. $x \in U_0$, where U_0 is as in theorem 2.1, and we are working on $SL'|_{q^{-1}(U_0)} = S^1 \times q^{-1}(U_0)$. So we can lift some homotopy (inside $q^{-1}(U_0)$) with boundary $\partial H'$ to one relating $\partial \tilde{H}$ to $(k')^n$. Hence, q'_* is an isomorphism.

Seifert proved that $\pi_1^V(SL) =$

$$\langle a_j, b_j, g_i, k : [a_j, k] = [b_j, k] = [g_i, k] = 1 = g_i^{m_i} k^{\beta_i} = k^{-b} \prod_{j=1}^g [a_j, b_j] \prod_{i=1}^n g_i \rangle$$

where g is the genus of Σ and m_i is the multiplicity of σ_i . Furthermore, the isotropy $\beta_i \bmod m_i$ of L at σ_i and $\deg L = b + \sum_1^n \beta_i / m_i$ are independent of the choices of lifts g_i, a_j and b_j of the generators of $\pi_1^V(\Sigma)$. By the isomorphism of the above extensions and if SL' is trivial, we can choose lifts such that all β_i 's are zero as well as b . By theorem 2.2, this implies that L is trivial. $\square$

Now we prove proposition 4.5:

Proof: Theorem 4.1 implies the existence of some $L \rightarrow \Sigma$ with $\det E' = q^*L$ and that the parabolic filtration of E' along P' is by trivial bundles; in particular $c_2(E') = 0$. Proposition 6.1 gives the uniqueness of L .

Now, $|E'|$ is uniquely determined by $\det E'$. Two different filtrations of $E'|_{P'_j}$ by trivial subbundles are related by a map $P'_j \rightarrow \mathrm{Sl}(r, \mathcal{C})$ which can be extended to an isomorphism of E' being the identity outside a tubular neighbourhood of P'_j since $\mathrm{Sl}(r, \mathcal{C})$ is simply connected. This shows (i). After the last argument and by theorem 2.2, part (ii) follows from proposition 6.1 and theorem 4.1. $\square$

References

- [AB] Atiyah, M. F., Bott, R.: The Yang-Mills equation over Riemann surfaces. Trans. R. Soc. Lond. A **308**, 523-615 (1982)

- [BPV] Barth, W., Peters, C., Van de Ven, A.: Compact Complex Surfaces. Berlin Heidelberg New York: Springer 1984
- [B] Bauer, S.: Parabolic bundles, elliptic surfaces and $SU(2)$ -representation spaces of genus zero Fuchsian groups. Math. Ann. **290**, 509-526 (1991)
- [BO] Bauer, S., Okonek, C.: The algebraic geometry of representation spaces associated to Seifert fibred homology 3-spheres. Math. Ann. **286**, 45-76 (1990)
- [B92] Biquard, O.: Prolongement d'un Fibre Holomorphe Hermitien a Courbure L^p sur une Courbe Ouverte. Int. J. Math. (3) **4**, 441-453 (1992)
- [B93] Biquard, O.: Sur les fibres paraboliques sur une surface complexe. preprint (1993)
- [Dol] Dolgachev. I.: Algebraic surfaces with $p_g = q = 0$. In: Algebraic surfaces CIME 1977, Liguori, Napoli, 97-215 (1981)
- [Don] Donaldson, S. K.: Anti-self-dual Yang-Mills connections over complex algebraic surfaces and stable vector bundles. Proc. London Math. Soc. (3) **50**, 1-26 (1985)
- [DK] Donaldson, S. K., Kronheimer, P. B.: The Geometry of Four-Manifolds. Oxford: Clarendon Press 1990
- [FS] Furuta, M., Steer, B.: Seifert Fibred Homology 3-Spheres and the Yang-Mills Equations on Riemann Surfaces with Marked Points. Adv. Math. (1) **96**, 38-102 (1992)
- [G] Gantz, C.: DPhil-Thesis, Oxford, in preparation
- [GH] Griffith, P., Harris, J.: Principles of Algebraic Geometry. New York: John Wiley & Sons 1978
- [KM93] Kronheimer, P. B., Mrowka, T. S.: Gauge Theory for Embedded Surfaces, I. Topology (4) **32**, 773-826 (1993)
- [KM95] Kronheimer, P. B., Mrowka, T. S.: Gauge Theory for Embedded Surfaces, II. Topology (1) **34**, 37-98 (1995)
- [L] Lübke, M.: Stability of Einstein-Hermitian Vector Bundles. Manus. Math. **42**, 245-257 (1983)

- [MS] Mehta, V. B., Seshadri, C. S.: Moduli of Vector Bundles on Curves with Parabolic Structures. *Math. Ann.* **248**, 205-239 (1980)
- [Mi] Miyaoka, Y.: Kähler metrics on elliptic surfaces. *Proc. Jap. Acad.* **50**, 533-536 (1974)
- [Mo] Moishezon, B.: Complex Surfaces and Connected Sums of Complex Projective Planes (Lect. Not. Math. 603) Berlin: Springer 1977
- [Mu] Munari, A.: Singular Instantons and Parabolic Bundles over Complex Surfaces (Oxford University, DPhil-Thesis) Oxford: 1993
- [NarS] Narasimhan, M. S., Seshadri, C. S.: Stable and Unitary Vector Bundles on Compact Riemann Surfaces. *Ann. Math.* **82**, 540-567 (1963)
- [NasS] Nasatyr, E. B., Steer, B.: The Narasimhan-Seshadri Theorem for Parabolic Bundles with Irrational Weights. preprint, Oxford (1993)
- [S] Simpson, C.: Constructing Variations of Hodge Structure using Yang-Mills-Theory. *J. Amer. Math. Soc.* **1**, 867-918 (1988)
- [SW] Steer, B., Wren, A.: The Donaldson-Hitchin-Kobayashi Correspondence for Parabolic Bundles over Orbifold Surfaces. preprint, Oxford (1994)
- [U] Ue, M.: On the Diffeomorphism Types of Elliptic Surfaces with Multiple Fibres. *Invent. Math.* **84**, 633-643 (1986)
- [UY] Uhlenbeck, K. K., Yau, S. T.: On the Existence of Hermitian-Yang-Mills Connections in Stable Vector Bundles. *Comm. Pure Appl. Math.* **39**, S257-S293 (1986)